

Upper bounds on r-mode amplitudes from observations of LMXB neutron stars

Simin Mahmoodifar¹ and Tod Strohmayer²

¹*Department of Physics, University of Maryland College Park, MD 20742, USA*

²*Astrophysics Science Division, NASA's Goddard Space Flight Center, Greenbelt, MD 20771, USA*

The r-mode oscillations of neutron stars can be potentially powerful probes of cold ultra-dense matter, since they couple the low energy degrees of freedom of dense neutron star matter to macroscopic, dynamic observables such as the stellar spin period and temperature. In this paper we present upper limits on the amplitude of r-mode oscillations, and their gravitational-radiation-induced spin-down rates, in low mass X-ray binary (LMXB) neutron stars under the assumption that the quiescent neutron star luminosity is powered by dissipation from a steady-state r-mode. We compute results for neutron star models constructed with the APR equation of state for masses of 1.4, 2 and 2.21 $M_{\odot}$. For the lower mass models (1.4 and 2 $M_{\odot}$) we find dimensionless r-mode amplitudes in the range from about 1×10^{-8} to 1.5×10^{-6} . For the accreting millisecond X-ray pulsar (AMXP) sources with known quiescent spin-down rates these limits suggest that $\lesssim 1\%$ of the observed rate can be due to an unstable r-mode. Interestingly, the AMXP with the highest amplitude limit, NGC 6640, could have an r-mode spin-down rate comparable to the observed, quiescent rate for SAX J1808-3658. Thus, quiescent spin-down measurements for this source would be particularly interesting. For all the sources considered here our amplitude limits suggest that their gravitational wave signals are likely too weak for detection with Advanced LIGO. Our highest mass model (2.21 $M_{\odot}$) can support enhanced, direct Urca neutrino emission in the core and thus can have higher r-mode amplitudes. Indeed, the inferred r-mode spin-down rates at these higher amplitudes are inconsistent with the observed spin-down rates for some of the sources, such as IGR J00291+5934 and XTE J1751-305. This can be used to place an upper limit on the masses of these sources if they are made of normal nuclear matter, or alternatively it could be used to probe the existence of exotic matter in them if their masses were known.

I. INTRODUCTION

The r-modes are large-scale currents in neutron stars that couple to gravitational radiation and remove energy and angular momentum from the star in the form of gravitational waves [1–4]. The physics of these oscillations is important in relating the microscopic properties of dense matter—such as its viscosity and emissivity—to the macroscopic, observable properties of neutron stars such as their spin frequency, temperature, mass and radius. The r-modes are unstable to gravitational radiation and their amplitudes can grow exponentially if viscous and other possible damping mechanisms are not large enough. Because they can affect the dynamic properties of the star—its spin frequency and temperature evolution—they are potentially important probes of the phases of dense matter. For example, the boundary of the r-mode instability region in the spin frequency - temperature plane, and in particular its minimum, which may determine the final rotation frequency of the star, is very different for stars with different interior compositions [5–7].

Whether or not the r-mode instability limits the spin rates of some neutron stars it is nevertheless important to explore for its potential astrophysically observable signatures. While mass-radius measurements of neutron stars are important for constraining the equation of state (EOS) of dense matter, observations of their dynamic properties such as spin and thermal evolution are important and more efficient in discriminating between different phases of dense matter. That's because dynamic properties are affected by the transport and thermody-

namic properties of dense matter inside the star, such as viscosity, heat conductivity and emissivity which depend on low energy degrees of freedom and are very different depending on the phase of dense matter present [5, 8, 9].

Pulse timing observations of accreting millisecond X-ray pulsars (AMXPs) made with NASA's Rossi X-ray Timing Explorer (RXTE) are beginning to reveal the long-term spin evolution of low mass X-ray binary (LMXB) neutron stars. The pulsar recycling hypothesis whereby millisecond radio pulsars acquire their fast spins via accretion, requires that at least some of these stars are spun-up to hundreds of Hz—the current neutron star spin record being 716 Hz [10]. Still, it remains somewhat puzzling that the spin frequency distribution of AMXPs appears to cut-off well below the mass-shedding limit of essentially all realistic neutron star EOS [11], and it has been suggested that the r-mode instability may play a role in limiting neutron star spin rates [12, 13]. Alternatively, recent work supports the idea that at least some of the AMXP population are in or close to “spin equilibrium,” essentially determined by the physics of magnetized accretion, and that an additional mechanism, such as the r-mode torque, is not required to halt the spin-up [14, 15].

While pulse timing noise in AMXPs has made interpretation of the X-ray pulsation data difficult, there are now several convincing measurements of spin-down during the quiescent phases between accretion outbursts. For example, both SAX J1808.4-3658 (hereafter SAX J1808) and IGR J00291+5934 (hereafter IGR J00291) show spin-down of the neutron star between outbursts, at rates, $\dot{\nu}$, of approximately $1 - 3 \times 10^{-15}$ Hz s⁻¹ [16, 17].

The most convincing evidence for an accretion-induced spin-up during outbursts is in IGR J00291, for which a peak value of about $3 \times 10^{-13} \text{ Hz s}^{-1}$ has now been inferred [16]. The magnitude of the “instantaneous” spin-up torque is larger in this source than the spin-down torque by about a factor of 100. The long-term evolution is then a competition between the accretion outbursts which spin it up and the quiescent spin-down intervals. If the outbursts are frequent enough then the star will be spun-up, if not, then long-term spin-down occurs. As the spin-up torque is proportional to the mass accretion rate, the outcome can also be expressed in terms of the long-term, average mass accretion rate.

The observed quiescent spin-down rate, $\dot{\nu}_{sd}$, puts an upper limit on the torque that can be present due to any r-mode, as it has to be less than the observed torque of $2\pi I \dot{\nu}_{sd}$, where I is the moment of inertia of the neutron star. Now, spin-down of pulsars is typically ascribed to the magnetic-dipole torque that is almost certainly present at some level due to the large scale magnetic field of the neutron star. Indeed, magnetic field estimates are usually obtained by equating the *entire* observed spin-down to that expected theoretically for magnetic-dipole radiation. But the observed spin-down is due to the sum total of any torques present. Assuming that both r-mode and magnetic-dipole torques are present, then the magnitude of the r-mode torque is equal to the observed torque minus the magnetic-dipole torque. Since the magnetic field strength and geometry of AMXPs are typically not known independently from spin-down estimates, the precise value of the magnetic-dipole torque is not known a priori. However, to the extent that the magnetic torque accounts for the majority of the observed spin-down, as is typically assumed, then the r-mode torque is likely to be much less than the measured spin-down torque.

In addition to torquing the neutron star the viscous damping of the r-modes acts as an internal source of heat. Calculations of the coupled thermal and spin evolution of accreting neutron stars including the effects of r-mode heating and gravitational radiation have been carried out by several authors [6, 7, 18, 19]. The primary assumption in these analyses has been that the long-term accretion (spin-up) torque is balanced by the r-mode torque due to the emission of gravitational radiation that carries away angular momentum. That is, the long-term average accretion torque is in equilibrium with the r-mode torque. As noted above, the recent pulse timing observations of AMXPs indicate that this equilibrium assumption is likely not realized in practice. Moreover, Brown & Ushomirsky [18] showed that if the average, fiducial accretion torque given by $N_{acc} = \langle \dot{M} \rangle (GMR)^{1/2}$, where $\langle \dot{M} \rangle$ is the long-term average mass accretion rate, were balanced by the r-mode torque then the quiescent luminosities of some accreting neutron star transients should be substantially larger than observed due to heating from the r-modes. Here we use a similar argument to place upper bounds on the r-mode amplitudes that can be present in accreting neutron stars assuming that r-mode heating

provides the source of neutron star luminosity in the absence of accretion. We then use these amplitude limits to assess the level of r-mode spin-down that can be present, its relation to observed spin-down rates when available, and the expected strength of gravitational radiation.

The paper is organized as follows. In §2 we review the basic theory of the r-mode instability and how it couples to the thermal and spin evolution of neutron stars. In §3 we describe our methods for constraining the r-mode amplitudes using the observed properties of quiescent, LMXB neutron stars, and we assess the implications for r-mode spin-down and the emission of gravitational radiation. We provide a summary and discussion of the implications of our findings in §4.

II. SPIN-DOWN DUE TO UNSTABLE R-MODES

The “flow pattern” of the r-modes is prograde in the inertial frame and retrograde in the rotating frame, which means that it moves in the same direction as the star’s rotation as seen by an observer at infinity, but in the opposite direction as seen by an observer at rest on the star. Any mode that is retrograde in the co-rotating frame and prograde in the inertial frame grows as a result of its emitting gravitational waves. This is the well-known Chandrasekhar-Friedman-Schutz (CFS) mechanism, and it means that gravitational radiation drives the mode rather than damps it. Viscosity on the other hand tends to damp the r-mode by transferring angular momentum from the mode to the rigid (unperturbed) star. The total angular momentum of a perturbed star can be written as:

$$J = I\Omega - J_c \quad (1)$$

where I is the moment of inertia and Ω is the angular rotation frequency of the star and J_c is the canonical angular momentum of the mode, given by:

$$J_c = -\frac{3}{2}\tilde{J}MR^2\Omega\alpha^2 \quad (2)$$

where α is the amplitude of the r-mode and $\tilde{J}$ is a dimensionless constant defined by [20]:

$$\tilde{J} \equiv \frac{1}{MR^4} \int_0^R \rho r^6 dr \quad (3)$$

where ρ is the run of density within the neutron star. The moment of inertia of the star, I , can also be written as $I = \tilde{I}MR^2$ where $\tilde{I}$ is defined by:

$$\tilde{I} \equiv \frac{8\pi}{3MR^2} \int_0^R \rho r^4 dr. \quad (4)$$

To derive the equations for the dynamical evolution of the star we use the following argument by Ho & Lai [21]:

The canonical angular momentum of the mode increases through gravitational radiation and decreases by

transferring angular momentum to the star through viscosity

$$\frac{dJ_c}{dt} = -\frac{2}{\tau_G}J_c - \frac{2}{\tau_V}J_c \quad (5)$$

where the viscous damping time τ_V is given by $\frac{1}{\tau_V} = \frac{1}{\tau_S} + \frac{1}{\tau_B} + \dots$. Here, τ_S and τ_B refer to shear and bulk viscosities, respectively, and the ellipsis denotes other possible dissipative mechanisms, such as boundary layer effects [22, 23].

The second evolution equation is obtained by writing the conservation of the total angular momentum J of the perturbed star which says that the total angular momentum of the star decreases due to gravitational radiation and increases due to accretion:

$$\frac{dJ}{dt} = -\frac{2J_c}{\tau_G} + N_{acc}, \quad (6)$$

where N_{acc} is the accretion torque. For a fiducial torque N_{acc} can be written as $\dot{M}(GMR)^{1/2}$ which assumes that each accreted particle transfers to the star an angular momentum equal to the Keplerian value at the stellar radius R [18]. In the previous equations the quantities $\frac{1}{\tau_i}$ are given by,

$$\frac{1}{\tau_i} = -\frac{P_i}{2E_c} \quad (7)$$

where E_c is the canonical energy of the r-mode, P_G is the power radiated by gravitational waves and P_B and P_S are the powers dissipated due to bulk and shear viscosity, respectively (in natural units where $c = \hbar = k_B = 1$),

$$E_c = \frac{1}{2}\alpha^2\Omega^2\tilde{J}MR^2 \quad (8)$$

$$P_G = \frac{32\pi(m-1)^{2m}(m+2)^{2m+2}}{((2m+1)!!)^2(m+1)^{2m+2}}\tilde{J}_m^2GM^2R^{2m+2}\alpha^2\Omega^{2m+4} \quad (9)$$

$$P_B = -\frac{16m}{(2m+3)(m+1)^5\kappa^2}\frac{\tilde{V}_m\Lambda_{\text{QCD}}^{9-\delta}R^7\alpha^2\Omega^4T^\delta}{\Lambda_{EW}^4} \quad (10)$$

$$P_S = -\frac{(m-1)(2m+1)\tilde{S}_m\Lambda_{\text{QCD}}^{3+\sigma}R^3\alpha^2\Omega^2}{T^\sigma}. \quad (11)$$

Here we consider $m = 2$ r-modes, and Λ_{QCD} and Λ_{EW} are characteristic strong and electroweak scales introduced to make $\tilde{V}$ and $\tilde{S}$ dimensionless. In our calculations we have used $\Lambda_{\text{QCD}} = 1$ GeV and $\Lambda_{EW} = 100$ GeV. The dimensionless parameters $\tilde{V}$, $\tilde{S}$, $\tilde{I}$ and $\tilde{J}$, which involve radial integration over the star, and δ and σ are given in Table I for three different neutron star models that we study in this paper. All three models here are made of normal hadronic matter with the APR equation of state [24] but they have different masses ($1.4 M_\odot$, $2.0 M_\odot$ and $2.21 M_\odot$) and radii [5]. The two models with

masses of $1.4 M_\odot$ and $2.0 M_\odot$ only allow modified Urca neutrino emission in the core, but the one with a mass of $2.21 M_\odot$ allows direct Urca neutrino emission in a core of radius 5.9 km. Direct Urca processes are very sensitive to the proton fraction of dense matter. The required proton fraction is roughly 14% in the case of the APR equation of state, reached at relatively high density $n \sim 5n_0$, where n_0 is the nuclear saturation density, this could be different for other equations of state [5].

The evolution equations for the amplitude of the r-mode, α , and spin frequency of the star, Ω , can be written by using equations 5 and 6 and substituting J_c from Eq. 2. The third equation –which describes the temperature evolution– can be obtained by noting that the temperature of the star decreases due to thermal emission from the surface and neutrino emission from the interior, which in an average mass hadronic star is dominated by modified Urca processes (in a massive star direct Urca processes can also happen in the core and should be included in the neutrino emissivity as well), and it increases due to the viscous dissipation of the r-mode energy, P_V . This gives the following equations for the evolution of spin frequency, r-mode amplitude and temperature,

$$\frac{d\Omega}{dt} = -2Q\frac{\Omega\alpha^2}{\tau_V} + \frac{N_{acc}}{I} \quad (12a)$$

$$\frac{d\alpha}{dt} = -\frac{\alpha}{\tau_G} - \frac{\alpha}{\tau_V}(1 - \alpha^2Q) - \frac{\alpha}{2\Omega}\frac{N_{acc}}{I} \quad (12b)$$

$$\frac{dE}{dt} = C_V\frac{dT}{dt} = -L_\nu - L_\gamma + |P_V|, \quad (12c)$$

where $Q \equiv \frac{3\tilde{J}}{2\tilde{I}}$, and the viscous dissipated power is $P_V = P_S + P_B + \dots$, and again the ellipsis denotes other possible dissipative processes.

Here, L_ν , L_γ and C_V , which are the total neutrino luminosity, photon luminosity and specific heat of the star, are given by (in natural units)

$$L_\nu = \frac{4\pi R^3\Lambda_{\text{QCD}}^{9-\theta}\tilde{L}}{\Lambda_{EW}^4}T^\theta \quad (13)$$

$$L_\gamma = 4\pi R^2\sigma T_{eff}^4 \quad (14)$$

$$C_V = 4\pi\Lambda_{\text{QCD}}^{3-v}R^3\tilde{C}_VT^v, \quad (15)$$

where T is the core temperature of the star and T_{eff} is the surface temperature. The dimensionless parameters $\tilde{L}_\nu$ and $\tilde{C}_V$ (that involve radial integration over the star) and θ and v are given in Table I for different stellar models.

Fig. 1 shows the r-mode instability window computed for a $1.4 M_\odot$ neutron star using the APR EOS (the modification to the instability window is rather modest in the case of the two other stellar models considered in this paper [5]). In the region above the curve the gravitational radiation timescale is smaller than the viscous damping timescales, therefore r-modes are unstable and their amplitudes grow exponentially in this region. As can be

Neutron Star	Shell	$R[km]$	$\Omega_K[Hz]$	$\tilde{I}$	$\tilde{J}$	$\tilde{S}$	$\tilde{V}$	$\tilde{C}_V$	$\tilde{L}$	σ	δ	v	θ
NS $1.4 M_\odot$	core	11.5	6020	0.283	1.81×10^{-2}	7.68×10^{-5}	1.31×10^{-3}	2.36×10^{-2}	1.91×10^{-2}	$\frac{5}{3}$	6	1	8
NS $2.0 M_\odot$		11.0	7670	0.300	2.05×10^{-2}	2.25×10^{-4}	1.16×10^{-3}	2.64×10^{-2}	1.69×10^{-2}				
NS $2.21 M_\odot$	m.U. core	10.0	9310	0.295	2.02×10^{-2}	5.05×10^{-4}	9.34×10^{-4}	2.62×10^{-2}	1.29×10^{-2}	$\frac{5}{3}$	4	1	6
	d.U. core	(5.9)					1.16×10^{-8}		2.31×10^{-5}				

Table I: Radius, Kepler frequency and radial integral parameters that appear in the moment of inertia, angular momentum of the mode, dissipative powers due to shear viscosity (from leptonic interactions) and bulk viscosity (due to Urca processes), specific heat and neutrino luminosity for different neutron star models considered in this work [5].

seen in this figure most of the LMXB's are in the unstable region for normal hadronic stars where the damping is due to the shear and bulk viscosity from leptonic scattering and Urca processes, therefore there must be some non-linear mechanism that saturates the amplitude of the unstable r-modes at a finite value. Supra-thermal bulk viscosity [25] is one of these non-linear mechanisms, but in the case that only the core of the star is considered and the effects of the crust are ignored, it can only saturate the r-mode at large amplitudes [26]. Mode coupling is another mechanism that can saturate the r-mode at small amplitude when hyperons and therefore large hyperon bulk viscosity is included in the computations [27, 28]. At this point it is not entirely clear which mechanism is actually responsible for saturating the r-mode amplitude, however, in this paper our primary interest is to obtain upper bounds on r-mode amplitudes from observations of neutron star transients and this does not require a precise understanding of the saturation mechanism. Understanding the detailed physics of the saturation mechanism is an important issue, but it is beyond the scope of this paper.

Here we note that if we were to consider the existence of exotic forms of matter in the star, then the shape of the instability window would change due to their different shear and bulk viscosities [5, 7, 29]. The shape of the instability window may also be different due to crust boundary layer effects [6], but even in that case for realistic models most of the LMXB sources are in the unstable region. Therefore, we assume that these sources have unstable r-modes which are emitting gravitational waves but the amplitude of the r-mode is not growing exponentially, and there is a mechanism that can saturate the growth of the r-mode. In addition, we assume that all of these stars are made of normal hadronic matter (constructed with the APR EOS) and all of the sources that are in the unstable region for hadronic stars in Fig. 1 are emitting gravitational waves at a constant amplitude.

Since we don't know which mechanism is actually responsible for saturating the r-mode amplitude, we simply assume that there is a nonlinear mechanism that saturates the mode. When α hits the saturation amplitude, the right hand side of Eq. 12b becomes zero which im-




Figure 1: The r-mode instability region for a $1.4M_\odot$ NS constructed with the APR equation of state in the spin frequency vs. core temperature plane. Also shown are some of the low mass X-ray binaries which have been considered in this work. The horizontal line extending rightward from the temperature symbols (the black squares) shows the difference between two models for relating the surface temperature to the core temperature (i.e. the difference between a fully or partially accreted envelope). The difference between the core temperatures in these two cases gets larger as the surface temperature increases, but even for the highest surface temperature considered in this work the difference is not big enough to change our results.

plies that

$$\frac{1}{\tau_V} = \frac{1}{\tau_G} \frac{1}{1 - \alpha^2 Q}, \quad (16)$$

where we have neglected the last term in Eq. 12b since it is much smaller than the other terms.

Therefore, when the amplitude is saturated, in all of the evolution equations $\frac{1}{\tau_V}$ should be replaced by $\frac{1}{\tau_G}(\frac{1}{1 - \alpha^2 Q})$ and $|P_V|$ by $\frac{P_G}{1 - \alpha^2 Q}$. Since viscosity alone can not stop the growth of the r-mode and an extra mechanism is needed to do that, therefore at saturation

$\frac{1}{\tau_G}(\frac{1}{1-\alpha^2 Q})$ is larger than $\frac{1}{\tau_V}$ and the reheating term on the right hand side of the temperature evolution equation (Eq 12c) will be larger than the reheating due to bulk and shear viscosity [8].

Source	$\Delta = \frac{t_o}{t_r}$	$\dot{\nu}_{su}$ (Hz s ⁻¹)	$\alpha_{sp.eq}$ (1.4M _⊙)	$\alpha_{sp.eq}$ (2.0M _⊙)	$\alpha_{sp.eq}$ (2.21M _⊙)
IGR J00291	$\frac{13}{1363}$	5×10^{-13}	1.46×10^{-7}	1.22×10^{-7}	1.41×10^{-7}
SAX J1808-3658	$\frac{40}{2 \times 365}$	2.5×10^{-14}	3.20×10^{-7}	2.66×10^{-7}	3.08×10^{-7}
XTE J1814-338	$\frac{50}{19 \times 365}$	1.5×10^{-14}	2.12×10^{-7}	1.76×10^{-7}	2.04×10^{-7}

Table II: Duty cycle, spin-up rate [30] and upper limits on r-mode amplitudes from the “spin-equilibrium” condition for three neutron star transients with assumed masses of 1.4M_⊙, 2.0M_⊙ and 2.21M_⊙.

III. CONSTRAINING THE R-MODE AMPLITUDE

A. Constraints from “spin equilibrium”

Here we compare two different methods for constraining the r-mode amplitude, α , from the observations of LMXB neutron star transients. The first one, which gives larger values for α , is based on the spin equilibrium assumption [6, 18, 19, 31] where we assume that in an outburst-quiescence cycle all the spin-up torque due to accretion during the outburst is balanced by the r-mode spin-down torque due to gravitational radiation in the whole cycle. This is similar to the prescription considered by previous authors ([6, 18]), but rather than using a “fiducial” torque estimated from the long-term average $\dot{M}$, we can now use the observed spin-up rates and outburst properties to directly constrain the torque.

Therefore we have:

$$2\pi I \dot{\nu} \Delta = \frac{2J_c}{\tau_G} \quad (17)$$

where $\dot{\nu}$ is the spin-up rate during outburst and $\Delta = \frac{t_o}{t_r}$ is the ratio of the outburst duration, t_o , to the recurrence time, t_r . We can estimate the left hand side of Eq. 17 from X-ray observations of LMXBs and since the right hand side is a function of α through J_c (see Eq. 2) we can determine α . In Table II we give the results for α for three sources for which there are now estimates of the spin-up rate due to accretion. The values of α are given for the three different neutron star models considered in this work, and they are all in the range from $\approx 1 - 3 \times 10^{-7}$.

At these amplitudes the inferred r-mode spin-down rate would be competitive with the magnetic dipole spin-down rate which almost certainly exists in these LMXBs, and which is quite likely the dominant spin-down mechanism. Moreover, the amplitudes are also comparable to those deduced assuming r-mode spin equilibrium with the “fiducial” accretion torques estimated by Brown & Ushomirsky [18]. These authors also demonstrated that

at such amplitudes some of these objects should have significantly higher quiescent luminosities due to r-mode reheating than observed. These results suggest that the r-mode torque does not balance the accretion-driven spin-up torque and that r-mode amplitude estimates based on the “spin equilibrium” assumption will overestimate the true amplitude.

B. Constraints from “thermal equilibrium”

Here, we use the same thermal equilibrium argument outlined by Brown & Ushomirsky, but rather than estimating the quiescent luminosity using the r-mode amplitude deduced from “spin equilibrium,” we use observations of the quiescent luminosity of LMXBs to directly constrain the amplitude of the r-mode. This works because in a steady-state, gravitational radiation pumps energy into the r-mode at a rate given by, $W_d = (1/3)\Omega \dot{J}_c = -2E_c/\tau_G$. This expression has the familiar relationship for a power dissipated by an applied torque, and in this case it is simply the r-mode torque due to gravitational radiation. In a thermal steady state all of this energy must be dissipated in the star. Some fraction of this heat will be lost from the star due to neutrino emission and the rest will be radiated at the surface. We further assume that all of the heat emitted from the surface of the star during quiescence is due to the r-mode dissipation inside the star. The resulting r-mode amplitude limits are upper bounds in the sense that the observed luminosity reflects the contribution from r-mode heating as well as any additional sources of heat that are present, such as for example due to accretion and the nuclear processing of accreted material. If any such sources of heat are present then the actual r-mode amplitude will be less than the upper bounds given here. For the sources that we study in this work, since we know the values of surface temperatures and quiescent luminosities from observations, we can estimate the core temperature and therefore determine the neutrino luminosities to estimate the total amount of heat deposited in the core of these systems by gravitational radiation.

To compute the core temperatures we use equation A8 in [35] which relates the effective surface temperature of the star T_{eff} to the internal temperature T_b , which is the temperature at a fiducial boundary at $\rho_b = 10^{10}$ g/cm³ for a fully accreted envelope and is valid at not too high temperatures ($T_b \leq 10^8$ K),

$$\left(\frac{T_{eff}}{10^6 K}\right)^4 = \left(\frac{g}{10^{14} cm s^{-2}}\right) (18.1 \frac{T_b}{10^9 K})^{2.42} \quad (18)$$

where $g = GM/(R^2 \sqrt{1 - r_g/R})$ is the surface gravity and $r_g = 2GM/c^2$. Here we assume that the neutron star’s core is isothermal and since the thermal conductivity of the crust is high [36] we have $T_{core} = T_b$ to good approximation. If we instead use equation A9 in [35] which gives T_b for a partially accreted envelope, and

Source	ν_s (Hz)	Distance (kpc)	kT_{eff} (eV)	$T_{core}(K)$ ($1.4M_\odot$)	$T_{core}(K)$ ($2.21M_\odot$)	$L_\gamma(erg.s^{-1})$ ($1.4M_\odot$)	$L_\nu(erg.s^{-1})$ ($1.4M_\odot$)	$L_\gamma(erg.s^{-1})$ ($2.21M_\odot$)	$L_\nu(erg.s^{-1})$ ($2.21M_\odot$)
4U 1608-522	620	4.1 ± 0.4	170	1.25×10^8	1.34×10^8	1.21×10^{34}	2.16×10^{32}	3.03×10^{34}	4.52×10^{38}
IGR J00291+5934	599	5 ± 1	71	2.96×10^7	3.18×10^7	3.65×10^{32}	2.06×10^{27}	9.27×10^{32}	7.94×10^{34}
MXB 1659-29	556	~ 10	55	1.96×10^7	2.07×10^7	1.35×10^{32}	7.64×10^{25}	3.29×10^{32}	6.08×10^{33}
Aql X-1	550	4.55 ± 1.35	94	4.70×10^7	5.06×10^7	1.12×10^{33}	8.40×10^{28}	2.87×10^{33}	1.31×10^{36}
KS 1731-260	524	7.2 ± 1.0	70	2.89×10^7	3.12×10^7	3.44×10^{32}	1.70×10^{27}	8.88×10^{32}	7.13×10^{34}
XTE J1751-305	435	9 ± 3	< 71	2.96×10^7	3.18×10^7	3.65×10^{32}	2.06×10^{27}	9.27×10^{32}	7.94×10^{34}
SAX J1808-3658	401	3.5 ± 0.1	< 30	7.23×10^6	7.69×10^6	1.21×10^{31}	2.63×10^{22}	2.99×10^{31}	1.60×10^{31}
XTE J1814-338	314	6.7 ± 2.9	< 69	2.82×10^7	3.01×10^7	3.24×10^{32}	1.39×10^{27}	8.12×10^{32}	5.72×10^{34}
NGC 6440	205	8.5 ± 0.4	87	4.11×10^7	4.46×10^7	8.10×10^{32}	2.88×10^{28}	2.11×10^{33}	6.10×10^{35}
XTE J1807-294	191	8.35 ± 3.65	< 51	1.72×10^7	1.83×10^7	9.84×10^{31}	2.71×10^{25}	2.46×10^{32}	2.95×10^{33}
XTE J0929-314	185	7.8 ± 4.2	< 50	1.66×10^7	1.79×10^7	9.06×10^{31}	2.06×10^{25}	2.31×10^{32}	2.53×10^{33}

Table III: Spin frequency, distance to the source [31], effective temperature at the surface of the star [32–34], core temperature, photon luminosity at the surface of the star and neutrino luminosity for both $1.4M_\odot$ & $2.2M_\odot$ NSs. Note that kT_{eff} given in this table is for a $1.4M_\odot$ NS with a radius of 10 km but in computing the core temperatures and luminosities for different neutron star models the appropriate red-shifts have been used. We note that for the sake of brevity the core temperatures for the $2M_\odot$ models are not included in the table, however, their values are all $\approx 5\%$ less than those for the $2.21M_\odot$ model.

a column depth of $\frac{P}{g} = 10^9$ g/cm² [7] we get core temperatures slightly higher than those derived from Eq. 18. The right hand side of the error bars on the temperatures in Fig.1 shows this difference. It is really only relevant for a single source, 4U 1608-522, but even in this case it is less than a 50% increase, and the difference is always small enough that it doesn't qualitatively change our results in the remainder of the paper. To compute T_b we have used the effective surface temperatures, T_{eff} , given by [32–34]¹. Note that those temperatures are computed for a $1.4M_\odot$ NS with a 10km radius and we have used the appropriate redshifts to compute T_{core} for our neutron star models.

Having surface and core temperatures for these sources we can use Eqs. 13 and 14 to evaluate their neutrino and thermal luminosities for different stellar models. The values of the core temperatures, photon and neutrino luminosities for different sources for $1.4M_\odot$ and $2.21M_\odot$ NS models are computed and given in Table III. Note that in the case of $1.4M_\odot$ NSs the neutrino luminosity is only due to the modified Urca reactions, but for the $2.21M_\odot$ NS it is due to both direct and modified Urca reactions. By comparing the neutrino and thermal luminosities in Table III one can see that at temperatures relevant for the LMXBs the standard neutrino cooling from modi-

fied Urca reactions is negligible compared to the photon emission from the surface of the star. However, if the star is massive enough to enable direct Urca reactions in the core (such as for our $2.21M_\odot$ neutron star model) then the neutrino emission will dominate the cooling process for surface temperatures higher than about 40 eV [38]. The thermal equilibrium condition can be written as $W_d = L_\nu + L_\gamma$, where reheating due to r-mode dissipation is given by $W_d = \frac{-2E_c}{\tau_{GR}}$ and is a function of r-mode amplitude, α . Therefore, α can be written in terms of luminosities as:

$$\alpha = \frac{5 \times 3^4}{2^8 \tilde{J} M R^3 \Omega^4} (\frac{L_\gamma + L_\nu}{2\pi G})^{1/2} \quad (19)$$

where $L_\gamma = 4\pi R^2 \sigma T_{eff}^4$ is the thermal photon luminosity at the surface of the star. Here, R and T_{eff} are the stellar radius and surface temperature, respectively, and the neutrino luminosity is given by

$$L_\nu = \frac{4\pi R_{DU}^3 \Lambda_{QCD}^3 \tilde{L}_{DU}}{\Lambda_{EW}^4} T^6 + \frac{4\pi(R^3 - R_{DU}^3) \Lambda_{QCD} \tilde{L}_{MU}}{\Lambda_{EW}^4} T^8 \quad (20)$$

where T is the core temperature, R_{DU} is the radius of the core where direct Urca neutrino emission is allowed and $\tilde{L}$ is a dimensionless parameter given in Table I. The thermal equilibrium condition for a NS with standard neutrino cooling ($1.4M_\odot$ and $2.0M_\odot$ NSs in this study) can be approximated as $W_d \simeq L_\gamma$, since the neutrino cooling in this case is negligible compared to the surface photon luminosity.

As can be seen in Fig.1, out of the 11 sources considered in this paper only 9 of them are likely to have unstable r-modes, meaning that they are above the r-mode

¹ It should be emphasized that the temperatures given in Table 2 of Heinke et. al. 2007 & 2009 [32, 33] are effective temperatures at the surface of the star and not redshifted surface temperatures seen by an observer at infinity. We note that they have been incorrectly assumed to be redshifted temperatures in references [7, 37].

Source	$\alpha_{th.eq}$ ($1.4M_{\odot}$)	$\alpha_{th.eq}$ ($2.0M_{\odot}$)	$\alpha_{th.eq}$ ($2.21M_{\odot}$)	$\dot{\nu}$ (Hz s $^{-1}$) ($1.4M_{\odot}$)	$\dot{\nu}$ (Hz s $^{-1}$) ($2.0M_{\odot}$)	$\dot{\nu}$ (Hz s $^{-1}$) ($2.21M_{\odot}$)	$\dot{\nu}_{sd}$ (Hz s $^{-1}$) observation
4U 1608-522	7.15×10^{-8}	6.60×10^{-8}	1.18×10^{-5}	-1.44×10^{-15}	-1.78×10^{-15}	-4.25×10^{-11}	
IGR J00291+5934	1.14×10^{-8}	1.32×10^{-8}	1.82×10^{-7}	-2.89×10^{-17}	-5.59×10^{-17}	-7.94×10^{-15}	-3×10^{-15}
MXB 1659-29	1.16×10^{-8}	1.07×10^{-8}	6.90×10^{-8}	-1.78×10^{-17}	-2.18×10^{-17}	-6.77×10^{-16}	
Aql X-1	3.49×10^{-8}	3.27×10^{-8}	1.02×10^{-6}	-1.49×10^{-16}	-1.89×10^{-16}	-1.37×10^{-13}	
KS 1731-260	2.35×10^{-8}	2.20×10^{-8}	2.93×10^{-7}	-4.81×10^{-17}	-6.09×10^{-17}	-8.07×10^{-15}	
XTE J1751-305	5.09×10^{-8}	4.76×10^{-8}	6.54×10^{-7}	-6.13×10^{-17}	-7.74×10^{-17}	-1.09×10^{-14}	-5.5×10^{-15}
SAX J1808-3658	1.28×10^{-8}	1.19×10^{-8}	2.16×10^{-8}	-2.19×10^{-18}	-2.74×10^{-18}	-6.74×10^{-18}	-5.5×10^{-16}
XTE J1814-338	1.76×10^{-7}	1.67×10^{-7}	2.04×10^{-6}	-7.49×10^{-17}	-9.73×10^{-17}	-1.08×10^{-14}	
NGC 6440	1.54×10^{-6}	1.45×10^{-6}	3.64×10^{-5}	-2.90×10^{-16}	-3.71×10^{-16}	-1.75×10^{-13}	

Table IV: Upper bounds on the r-mode amplitude from the “thermal equilibrium” condition that are consistent with quiescent luminosity data are given for different star models. The gravitational radiation induced spin-down rates due to unstable r-modes as well as the observed spin-down rate for some of the sources are also given [16, 30, 39].

instability curve. The two most slowly rotating sources, XTE J1807 and XTE J0929, are outside the instability region for a hadronic star, which means they likely can no longer spin down due to gravitational radiation from an r-mode. Therefore, we only evaluate the upper bounds on the r-mode amplitude for those 9 sources within the instability window. Using Eq. 19, we have evaluated α for all of those sources using the three different NS models considered in this work. The values of α are given in Table IV. As can be seen for the $1.4M_{\odot}$ and $2.0M_{\odot}$ NSs, where there is no enhanced neutrino emission, the values of α range from 1.14×10^{-8} to 1.54×10^{-6} , where NGC 6440—which has the lowest spin frequency—has the highest r-mode amplitude, as expected. In the case of $2.21M_{\odot}$ NSs the upper bounds on α are larger since due to the direct Urca neutrino emission W_d can be larger, and α in this case ranges from 2.16×10^{-8} to 3.64×10^{-5} . 4U 1608-522 has the highest temperature among the sources considered which implies a very large neutrino luminosity for the $2.21M_{\odot}$ NS model and as a result a large r-mode amplitude. But it should be noted that this system has been accreting for a long time and both its high quiescent luminosity and surface temperature may be due to the long term accretion and our assumption in ascribing all the quiescent luminosity to the heat that comes from inside of the star is likely not a good estimate for this system, but here we are only interested in obtaining upper limits on r-mode amplitude.

C. r-mode spin-down

To see whether these results are plausible or not and what fraction of the quiescent spin-down of these sources can be due to gravitational radiation from r-mode oscillations, we use our results for α from the “thermal equilibrium” condition and insert them into the right hand side of Eq. 12a to determine $\dot{\nu}$ in the absence of accretion

($N_{acc} = 0$). The derived spin-down values for different neutron star models are given in Table IV, and are shown graphically in Figure 2 (vertical green lines). Comparing these results with the observed spin-down rates, which exist for IGR J00291, XTE 1751-305 and SAX J1808 (red diamonds in Figure 2), we find that in the case of $1.4M_{\odot}$ (“x” symbols) and $2.0M_{\odot}$ (square symbols) NSs, where there is no fast neutrino cooling present in the star, the r-mode spin-down can only provide about 1% of the observed spin-down rate, which means that other spin-down mechanisms such as magnetic-dipole radiation are responsible for spinning down a $1.4M_{\odot}$ or $2.0M_{\odot}$ hadronic star with no fast cooling process. For the two AMXPs with relatively “slow” spin frequencies, NGC 6640 and XTE J1814-338, we find r-mode spin-down limits that are more competitive with observed values. Indeed, our limit for NGC 6640 is comparable to the measured, quiescent spin-down rate for SAX J1808, thus spin-down measurements for this source would be particularly interesting.

On the other hand, the values of spin-down rate that we get for a $2.21M_{\odot}$ (circle symbols) hadronic star with direct Urca neutrino emission is not consistent with the observations for most of the sources since it gives a very large rate and in the case of IGR J00291 and XTE J1814 it is larger than the observed values. However, it is consistent for SAX J1808 and perhaps MXB 1659-29, and in the case of SAX J1808 r-mode spin-down is still about 1% of the observed spin-down rate.

With typical values of a few $\times 10^{-8}$, our derived amplitude upper limits suggest that for many LMXB neutron stars the r-modes are likely not excited to sufficient amplitudes to substantially affect their spin evolution. This begs the question of whether or not unstable, steady-state r-modes actually exist in these neutron stars? One possibility is that additional damping mechanisms, such as perhaps associated with crust effects, superfluid mutual friction [6] or the existence of




Figure 2: Limits on the spin-down rates due to an r-mode torque for nine LMXB systems and a range of neutron star masses are shown in the $\dot{\nu}$ versus ν plane. The vertical green lines show the full range of $\dot{\nu}$ for each source (labelled) obtained from the r-mode amplitude limits derived from observations of quiescent luminosities and temperatures (see the discussion in §III). The limits obtained for 1.4 , 2.0 , and $2.21M_{\odot}$ NS models are marked by the “x,” square, and circle symbols, respectively. Also shown are two pairs of parallel lines representing magnetic-dipole (solid) and r-mode (dashed) braking laws. Lines are drawn for two values of the magnetic field, 10^8 G (lower), and 10^9 G (upper), as well as two values of α , 5×10^{-8} (lower) and 5×10^{-6} (upper). For systems with measured, quiescent spin-down rates, these values are marked with the red diamond symbols. For additional context, millisecond pulsars from the ATNF pulsar database [40] are shown and denoted by the black “+” symbols.

exotic matter in the core of NSs [5, 29], are at work and modify the instability window so as to render these neutron stars stable to r-mode excitation. An interesting, related question is whether the existence of r-modes at the amplitudes estimated here can be inferred directly from observations? Figure 2 shows both r-mode (dashed parallel lines) and magnetic-dipole (solid parallel lines) spin-down laws. The r-mode spin-down braking index, $n = 7$, is steeper compared to that for magnetic spin-down ($n = 3$), thus at high enough spin frequencies (well above a kHz) one might expect that the r-mode torque would eventually become competitive with or dominate the magnetic dipole torque. However, as of yet there are no known neutron stars spinning fast enough for this effect to become dominant, and depending on the EOS the mass-shedding limit might be reached before the r-mode torque becomes competitive with the magnetic torque.

As discussed above, quiescent spin-down measure-

ments have been typically attributed to the magnetic torque. For a number of sources considered here our r-mode amplitude limits support this presumption. Any spin-down contribution from an r-mode torque would be more easily identifiable if the magnetic field strengths of these neutron stars were constrained independently from the magnetic spin-down estimate. Moreover, identifying an r-mode spin-down would, in principle, be simpler for those neutron stars with the lowest magnetic torques, and thus field strengths. At present, the lowest inferred millisecond pulsar magnetic field strengths are $\approx 6 \times 10^7$ G. At this level the magnetic spin-down is of order $\dot{\nu} \approx 5 \times 10^{-17}$ Hz s $^{-1}$, which is comparable to our derived r-mode spin-down limits for a number of sources considered here, assuming their neutron stars are $< 2M_{\odot}$. In this regard, quiescent spin-down measurements for more of the sources considered here, in particular the AMXPs XTE J1814 and NGC 6440, would

be extremely valuable.

D. Gravitational wave amplitudes

R-modes in NSs are one of the possible mechanisms for gravitational wave (GW) emission and they can be observationally interesting in newborn neutron stars and perhaps accreting neutron stars in LMXBs [41]. Continuous GW emission from r-modes is dominated by $l = m = 2$ current quadrupole emission [4]. The gravitational wave amplitude h_0 (strain tensor amplitude) is related to the r-mode amplitude α by the following equation [41, 42]:

$$h_0 = \sqrt{\frac{8\pi}{5}} \frac{G}{c^5} \frac{1}{r} \alpha \omega_r^3 M R^3 \tilde{J}, \quad (21)$$

where r is the distance to the source, M and R are the mass and radius of the NS, $\tilde{J}$ is the dimensionless parameter defined by Eq. 3 and ω_r is the frequency of the r-mode, which is related to the angular spin frequency of the star Ω (for the $m = l = 2$ r-mode) by the following equation

$$\omega_r \approx \frac{4}{3} \Omega. \quad (22)$$

Using the upper limits on the r-mode amplitude of NSs in LMXBs derived above, we can obtain upper limits on the amplitude of the GWs emitted from these sources due to unstable r-modes. For the sources considered in this work, upper limits on the GW strain amplitude h_0 for the $1.4M_\odot$ and $2.0M_\odot$ NS models are in the range of 1.8×10^{-29} to 4.9×10^{-28} which is below the anticipated detectability threshold of Advanced LIGO [31, 43]. In the case of the $2.21M_\odot$ NS model, since the r-mode amplitudes are larger than those for the low mass NSs, we get larger values of h_0 , but even in this case for most of the sources h_0 is still below the detectability threshold of Advanced LIGO. The highest values of h_0 for a massive star are obtained for NGC 6440 and Aql X-1 with an amplitude of the order of 4×10^{-27} and 4U 1608-522 with an amplitude of 7.18×10^{-26} . However, it should be noted that the large values of r-mode amplitude in these sources, which would cause very large spin-down rates, may eventually be ruled out by future measurements of their spin-down rates. In this context it is important to restore the X-ray timing capability that was lost when RXTE was decommissioned in January 2012. Missions planned or currently in development which could provide such a capability include India's *ASTROSAT* [44], ESA's Large Observatory for X-ray Timing (LOFT, [45]), and NASA's Neutron Star Interior Composition Explorer (NICER, [46]).

The upper limits on GW amplitudes discussed here are related to the GW emission due to unstable r-modes but our results do not exclude the possibility of having larger GW amplitudes in LMXBs from other GW emission mechanisms such as NS mountains [47, 48]. It is

worth mentioning that indirect upper limits on GW amplitude can be obtained for sources with observed spin-down rates, $\dot{\nu}$, by assuming that all of the observed spin-down is due to GW emission [41]:

$$h_0^{sd} = \frac{1}{r} \sqrt{\frac{45GI\dot{P}}{8c^3P}} \quad (23)$$

where $P = \frac{2\pi}{\Omega}$ is the observed pulse period and $|\frac{\dot{P}}{P}| = |\frac{\dot{\nu}}{\nu}|$. Using this equation for the three sources with measured $\dot{\nu}$ and different NS models (i.e. different masses and radii) we get h_0^{sd} values that range from 4.14×10^{-28} to 6.53×10^{-28} .

IV. CONCLUSIONS

In this paper we have presented upper limits on r-mode amplitudes in LMXB neutron stars using their observed quiescent luminosities, temperatures and spin-down rates. We calculated results for NS models constructed with the APR EOS (normal hadronic matter) with masses of 1.4, 2 and $2.21 M_\odot$, where our highest mass model ($2.21 M_\odot$) can support enhanced, direct Urca neutrino emission in the core. We have used two different methods to calculate r-mode amplitudes. The first is based on the assumption that in an outburst-quiescence cycle all the spin-up torque due to accretion during the outburst is balanced by the r-mode spin-down torque due to gravitational radiation. This method gives amplitudes in the range of $\approx 1 - 3 \times 10^{-7}$ for the sources with measured spin-up rates. Since in reality there are other sources of spin-down such as magnetic-dipole radiation that may be the dominant spin-down source, we use another method for computing the r-mode amplitude that doesn't ascribe all of the spin-down of the star to gravitational radiation and therefore gives tighter bounds on the amplitudes. This second method is based on the assumption that in a thermal steady state some fraction of the heat that is generated in the star due to r-mode dissipation will be lost from the star by neutrino emission and the rest will be radiated at the surface. This assumes that all of the heat emitted from the surface of the star during quiescence is due to the r-mode dissipation inside the star, and thus provides an upper bound on the r-mode amplitude. We have computed core temperatures as well as neutrino and thermal (photon) luminosities for LMXB sources using measurements of the quiescent luminosities and surface temperatures and showed that at temperatures relevant for LMXB neutron stars, when there is no enhanced cooling mechanism, the cooling of the star is dominated by photon emission from the surface (for 1.4 and $2.0M_\odot$ NS models), but in a massive star where direct Urca neutrino emission is allowed, the cooling is dominated by neutrino emission (for $T_{eff} \gtrsim 40$ eV). For the lower mass NS models (1.4 and $2 M_\odot$) we find dimensionless r-mode amplitudes in the range from about 1×10^{-8} to 1.5×10^{-6} . For the AMXP sources

with known quiescent spin-down rates these limits suggest that $\lesssim 1\%$ of the observed rate can be due to an unstable r-mode. Interestingly, the AMXP with the highest amplitude limit, NGC 6640, could have an r-mode spin-down rate comparable to the observed, quiescent rate for SAX J1808. Thus, quiescent spin-down measurements for this source would be particularly interesting. Having enhanced, direct Urca neutrino emission in the core of our highest mass model ($2.21 M_{\odot}$) means that the dissipated heat in the star can be larger and therefore it can have higher r-mode amplitudes. Indeed, the inferred r-mode spin-down rates at these higher amplitudes are inconsistent with the observed spin-down rates for some of the LMXB sources, such as IGR J00291 and XTE J1751-305. This could be used to put an upper limit on the masses of these sources if they were made of normal nuclear matter, or it could be used to probe the existence of exotic matter in them if the NS mass in these systems were known. Using the results for r-mode amplitudes, the upper limits on gravitational wave amplitude due to r-modes have

been computed. The upper limits on the GW strain amplitude h_0 for the $1.4M_{\odot}$ and $2.0M_{\odot}$ NS models are in the range of 1.8×10^{-29} to 4.9×10^{-28} which is below the anticipated detectability threshold of Advanced LIGO. In the case of the $2.21M_{\odot}$ NS model we get larger values for h_0 , but even in this case for most of the sources considered in this work, h_0 is still below the detectability threshold of Advanced LIGO. Gravitational waves due to other mechanisms such as NS mountains may have larger amplitudes in these systems.

Acknowledgments

SM thanks Mark Alford, Andrew Cumming, Cole Miller and Kai Schwenzer for helpful discussions. TS acknowledges NASA's support for high energy astrophysics. SM acknowledges the support of the U.S. Department of Energy through grant number DEFG02-93ER-40762.

-
- [1] N. Andersson, *A new class of unstable modes of rotating relativistic stars*, *Astrophys. J.* **502** (1998) 708–713, [[gr-qc/9706075](#)].
 - [2] J. L. Friedman and S. M. Morsink, *Axial instability of rotating relativistic stars*, *Astrophys. J.* **502** (1998) 714–720, [[gr-qc/9706073](#)].
 - [3] J. L. Friedman and B. F. Schutz, *Secular instability of rotating Newtonian stars*, *Astrophys. J.* **222** (1978) 281.
 - [4] L. Lindblom, B. J. Owen, and S. M. Morsink, *Gravitational radiation instability in hot young neutron stars*, *Phys. Rev. Lett.* **80** (1998) 4843–4846, [[gr-qc/9803053](#)].
 - [5] M. Alford, S. Mahmoodifar, and K. Schwenzer, *Viscous damping of r-modes: Small amplitude instability*, *Phys. Rev. D* **85** (2012) 024007, [[arXiv:1012.4883](#)].
 - [6] W. C. Ho, N. Andersson, and B. Haskell, *Revealing the physics of r-modes in low-mass X-ray binaries*, *Phys. Rev. Lett.* **107** (2011) 101101, [[arXiv:1107.5064](#)].
 - [7] B. Haskell, N. Degenaar, and W. C. G. Ho, *Constraining the physics of the r-mode instability in neutron stars with X-ray and ultraviolet observations*, *Mon. Not. R. Astron. Soc.* **424** (July, 2012) 93–103, [[arXiv:1201.2101](#)].
 - [8] M. G. Alford, S. Mahmoodifar, and K. Schwenzer, *Impact of r-modes on the cooling of neutron stars*, *AIP Conf. Proc.* **1492** (2012) 257–263, [[arXiv:1209.5962](#)].
 - [9] M. G. Alford and K. Schwenzer, *Gravitational wave emission and spindown of young pulsars*, [arXiv:1210.6091](#).
 - [10] J. W. T. Hessels, S. M. Ransom, I. H. Stairs, P. C. C. Freire, V. M. Kaspi, and F. Camilo, *A Radio Pulsar Spinning at 716 Hz*, *Science* **311** (Mar., 2006) 1901–1904, [[astro-ph/](#)].
 - [11] D. Chakrabarty, E. H. Morgan, M. P. Muno, D. K. Galloway, R. Wijnands, M. van der Klis, and C. B. Markwardt, *Nuclear-powered millisecond pulsars and the maximum spin frequency of neutron stars*, *Nature (London)* **424** (July, 2003) 42–44, [[astro-ph/](#)].
 - [12] L. Bildsten, *Gravitational Radiation and Rotation of Accreting Neutron Stars*, *Astrophys. J. Lett.* **501** (July, 1998) L89, [[astro-ph/](#)].
 - [13] N. Andersson, K. D. Kokkotas, and N. Stergioulas, *On the relevance of the r mode instability for accreting neutron stars and white dwarfs*, *Astrophys. J.* **516** (1999) 307, [[astro-ph/9806089](#)].
 - [14] N. E. White and W. Zhang, *Millisecond X-Ray Pulsars in Low-mass X-Ray Binaries*, *Astrophys. J. Lett.* **490** (Nov., 1997) L87.
 - [15] A. Patruno, B. Haskell, and C. D'Angelo, *Gravitational Waves and the Maximum Spin Frequency of Neutron Stars*, *Astrophys. J.* **746** (Feb., 2012) 9, [[arXiv:1109.0536](#)].
 - [16] A. Patruno, *The Accreting Millisecond X-ray Pulsar IGR J00291+5934: Evidence for a Long Timescale Spin Evolution*, *Astrophys. J.* **722** (2010) 909–918, [[arXiv:1006.0815](#)].
 - [17] J. M. Hartman, A. Patruno, D. Chakrabarty, C. B. Markwardt, E. H. Morgan, M. van der Klis, and R. Wijnands, *A Decade of Timing an Accretion-powered Millisecond Pulsar: The Continuing Spin Down and Orbital Evolution of SAX J1808.4-3658*, *Astrophys. J.* **702** (Sept., 2009) 1673–1678, [[arXiv:0902.2112](#)].
 - [18] E. F. Brown and G. Ushomirsky, *Constraints on the steady-state r-mode amplitude in neutron star transients*, [astro-ph/9912113](#).
 - [19] B. Haskell and A. Patruno, *Spin equilibrium with or without gravitational wave emission: the case of XTE J1814-338 and SAX J1808.4-3658*, [arXiv:1106.6264](#).
 - [20] B. J. Owen *et. al.*, *Gravitational waves from hot young rapidly rotating neutron stars*, *Phys. Rev. D* **58** (1998) 084020, [[gr-qc/9804044](#)].
 - [21] W. C. Ho and D. Lai, *R mode oscillations and spindown of young rotating magnetic neutron stars*, *Astrophys. J.* **543** (2000) 386–394, [[astro-ph/9912296](#)].
 - [22] Y. Wu, C. D. Matzner, and P. Arras, *R modes in neutron stars with crusts: Turbulent saturation*,

- spindown, and crust melting*, *Astrophys.J.* **549** (2001) 1011–1020, [[astro-ph/0006123](#)].
- [23] L. Bildsten and G. Ushomirsky, *Viscous Boundary-Layer Damping of R-Modes in Neutron Stars*, *Astrophys. J. Lett.* **529** (Jan., 2000) L33–L36.
- [24] A. Akmal, V. R. Pandharipande, and D. G. Ravenhall, *The equation of state for nucleon matter and neutron star structure*, *Phys. Rev.* **C58** (1998) 1804–1828, [[nucl-th/9804027](#)].
- [25] M. G. Alford, S. Mahmoodifar, and K. Schwenzer, *Large amplitude behavior of the bulk viscosity of dense matter*, *J. Phys.* **G37** (2010) 125202, [[arXiv:1005.3769](#)].
- [26] M. G. Alford, S. Mahmoodifar, and K. Schwenzer, *Viscous damping of r-modes: Large amplitude saturation*, *Phys.Rev.* **D85** (2012) 044051, [[arXiv:1103.3521](#)].
- [27] P. Arras, E. E. Flanagan, S. M. Morsink, A. K. Schenk, S. A. Teukolsky, *et. al.*, *Saturation of the R mode instability*, *Astrophys.J.* **591** (2003) 1129–1151, [[astro-ph/0202345](#)].
- [28] R. Bondarescu, S. A. Teukolsky, and I. Wasserman, *Spinning down newborn neutron stars: nonlinear development of the r-mode instability*, *Phys.Rev.* **D79** (2009) 104003, [[arXiv:0809.3448](#)].
- [29] K. Schwenzer, *How long-range interactions tune the damping in compact stars*, [arXiv:1212.5242](#).
- [30] A. Patruno and A. Watts, *Accreting Millisecond X-Ray Pulsars*, [arXiv:1206.2727](#).
- [31] A. Watts, B. Krishnan, L. Bildsten, and B. F. Schutz, *Detecting gravitational wave emission from the known accreting neutron stars*, *Mon.Not.Roy.Astron.Soc.* **389** (2008) 839–868, [[arXiv:0803.4097](#)].
- [32] C. O. Heinke, P. Jonker, R. Wijnands, and R. Taam, *Constraints on Thermal X-ray Radiation from SAX J1808.4-3658 and Implications for Neutron Star Neutrino Emission*, *Astrophys.J.* **660** (2007) 1424–1427, [[astro-ph/0612232](#)].
- [33] C. Heinke, P. Jonker, R. Wijnands, C. Deloye, and R. Taam, *Further Constraints on Thermal Quiescent X-ray Emission from SAX J1808.4-3658*, *Astrophys.J.* **691** (2009) 1035–1041, [[arXiv:0810.0497](#)].
- [34] J. A. Tomsick, D. M. Gelino, J. P. Halpern, and P. Kaaret, *The Low quiescent x-ray luminosity of the neutron star transient XTE J2123-058*, *Astrophys.J.* **610** (2004) 933–940, [[astro-ph/0404287](#)].
- [35] A. Potekhin, G. Chabrier, and D. Yakovlev, *Internal temperatures and cooling of neutron stars with accreted envelopes*, [astro-ph/9706148](#).
- [36] E. F. Brown and A. Cumming, *Mapping crustal heating with the cooling lightcurves of quasi-persistent transients*, *Astrophys.J.* **698** (2009) 1020–1032, [[arXiv:0901.3115](#)].
- [37] N. Degenaar, A. Patruno, and R. Wijnands, *The quiescent X-ray properties of the accreting millisecond X-ray pulsar and eclipsing binary Swift J1749.4-2807*, *Astrophys.J.* **756** (2012) 148, [[arXiv:1204.6059](#)].
- [38] E. F. Brown, L. Bildsten, and R. E. Rutledge, *Crustal heating and quiescent emission from transiently accreting neutron stars*, [astro-ph/9807179](#).
- [39] A. Patruno, P. Bult, A. Gopakumar, J. Hartman, R. Wijnands, *et. al.*, *Accelerated Orbital Expansion And Secular Spin Down of the Accreting Millisecond Pulsar SAX J1808.4-3658*, [arXiv:1111.6967](#).
- [40] R. N. Manchester, G. B. Hobbs, A. Teoh, and M. Hobbs, *The Australia Telescope National Facility pulsar catalogue*, *Astron.J.* **129** (2005) 1993, [[astro-ph/0412641](#)].
- [41] B. J. Owen, *How to adapt broad-band gravitational-wave searches for r-modes*, *Phys.Rev.* **D82** (2010) 104002, [[arXiv:1006.1994](#)].
- [42] B. J. Owen, *How photon astronomy affects searches for continuous gravitational waves*, *Class.Quant.Grav.* **26** (2009) 204014, [[arXiv:0904.4848](#)].
- [43] **Virgo Collaboration** Collaboration, B. Abbott *et. al.*, *Searches for gravitational waves from known pulsars with S5 LIGO data*, *Astrophys.J.* **713** (2010) 671–685, [[arXiv:0909.3583](#)].
- [44] G. Stewart, *ASTROSAT: A Multi- Wavelength Satellite. 1st Dedicated Indian Astronomical Mission.*, in *Half a Century of X-Ray Astronomy, Proceedings of the conference held 17-21 September, 2012 on Mykonos Island, Greece. Online at http://www.astro.noa.gr/xcosmo id.95, Sept., 2012*.
- [45] M. Feroci, J. W. den Herder, E. Bozzo, D. Barret, S. Brandt, M. Hernanz, M. van der Klis, M. Pohl, A. Santangelo, L. Stella, and *et al.*, *LOFT: the Large Observatory For X-ray Timing*, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 8443 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Sept., 2012. [arXiv:1209.1497](#).
- [46] K. C. Gendreau, Z. Arzoumanian, and T. Okajima, *The Neutron star Interior Composition ExploreR (NICER): an Explorer mission of opportunity for soft x-ray timing spectroscopy*, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 8443 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Sept., 2012.
- [47] G. Ushomirsky, C. Cutler, and L. Bildsten, *Deformations of accreting neutron star crusts and gravitational wave emission*, *Mon.Not.Roy.Astron.Soc.* **319** (2000) 902, [[astro-ph/0001136](#)].
- [48] B. Haskell, D. Jones, and N. Andersson, *Mountains on Neutron Stars: Accreted vs. Non-Accreted crusts*, *Mon.Not.Roy.Astron.Soc.* **373** (2006) 1423–1439, [[astro-ph/0609438](#)].